

Salix bebbiana Shrubland

COMMON NAME	Gray Willow Shrubland
SYNONYM	Beaked Willow Scrub
PHYSIOGNOMIC CLASS	Shrubland (III)
PHYSIOGNOMIC SUBCLASS	Deciduous shrubland (III.B)
PHYSIOGNOMIC GROUP	Cold-deciduous shrubland (III.B.2)
PHYSIOGNOMIC SUBGROUP	Natural/Semi-natural (III.B.2.N)
FORMATION	Temporarily flooded cold-deciduous shrubland (III.B.2.N.d)
ALLIANCE	<i>Salix bebbiana</i> Temporarily Flooded Shrubland Alliance

CLASSIFICATION CONFIDENCE LEVEL 2

USFWS WETLAND SYSTEM Palustrine

RANGE

Globally

This community has been identified in the Black Hills of South Dakota and Wyoming. It also appears in Montana.

Wind Cave National Park

Bebb's willow shrubland does not occur in Wind Cave NP. It is found along streams in the area west of the Park.

ENVIRONMENTAL DESCRIPTION

Globally

This community is a briefly flooded scrub-shrub wetland on slightly to moderately alkaline soils, usually near low-gradient streams. The water table is well below the soil surface for over half the growing season. However, there are brief periods of several days to a few weeks when water is at the surface.

Wind Cave National Park

Bebb's willow shrubland occurs in narrow zones along streams in the Wind Cave NP area.

MOST ABUNDANT SPECIES

Globally

<u>Stratum</u>	<u>Species</u>
Shrub	<i>Salix bebbiana</i>
Herbaceous	<i>Carex</i> spp., <i>Scirpus</i> spp.

Wind Cave National Park

<u>Stratum</u>	<u>Species</u>
Shrub	<i>Salix bebbiana</i>

CHARACTERISTIC SPECIES

Globally

Salix bebbiana

Wind Cave National Park

Salix bebbiana

VEGETATION DESCRIPTION

Globally

This shrubland is typically dominated by a dense growth of shrubs 0.5-3 m tall. The most abundant species in the shrub layer are *Salix bebbiana*, *Salix scouleriana*, and *Salix fendleriana*. Other species found in this strata include *Betula occidentalis*, *Cornus sericea* ssp. *sericea*, *Salix exigua*, *Salix fluviatilis*, and *Prunus virginiana*. Multiple-stemmed trees and/or shrubs 1.5-5 m tall have 30-100% cover; single-stemmed trees have less than 30% cover. The herbaceous layer often contains *Scirpus* spp., *Carex* spp., *Triglochin palustris*, *Calamagrostis canadensis*, and *Equisetum* spp. along the wetter margins of the community. In the drier areas *Gentiana strictiflora*, *Prunella vulgaris*, *Pyrola asarifolia*, *Zizia cordata*, *Sanicula marilandica*, *Viola canadensis*, *Vicia americana*, and *Ranunculus macounii* are frequently present.

Wind Cave National Park

Bebb's Willow Shrubland is strongly dominated by Bebb's willow, with cover in the 50-75% range and sometimes greater. In the Wind Cave area, other shrubs that contribute minor cover are juneberry (*Amelanchier alnifolia*) and several species of currant (*Ribes* spp.). Herbaceous cover is sparse overall, but patches of western Great Plains streamside vegetation can be found in openings in the willows.

OTHER NOTEWORTHY SPECIES

CONSERVATION RANK G3?. Most sites have been impacted by grazing.

DATABASE CODE CEGL001173

MAP UNITS

The Bebb's willow community corresponds to map unit 34, Bebb Willow Shrubland, on the Wind Cave vegetation map.

COMMENTS

REFERENCES

Baker, W.L. 1982b. Natural vegetation of the Piceance Basin, Colorado. Appendix D, Pages 1-113 in Peterson, J.S. and W.L. Baker, eds., Inventory of the Piceance Basin, Colorado. Unpublished report done for the Bureau Land Management, Craig, CO.

Dick-Peddie, W.A. 1993. New Mexico Vegetation, Past, Present and Future. University of New Mexico Press. Albuquerque, NM. 244 PP.

Hansen, P., K. Boggs, R. Pfister. 1991. Classification and management of riparian and wetland sites in Montana. Unpublished draft version prepared for Montana Riparian Association, Montana Forest and Conservation Experiment Station, School of Forestry, University of Montana, Missoula, MT. 478 pp.

Hayward, H.E. 1928. Studies of plants in the Black Hills of South Dakota. Botanical Gazette 85(4):353-412.

Osborn, R., G. Kittel, and M. Reid. 1998. Colorado Riparian Plant Associations and Western States Vegetation Classification. CDROM. U.S. Geological Survey, Mid-Continent Ecology Research Center, Fort Collins, CO.

Szaro, R.C. 1989. Riparian forest and scrubland community types of Arizona and New Mexico. Desert Plants Special Issue 9(3-4):70-139.

The Nature Conservancy (TNC). 1991. Community characterization abstract - alkaline scrub riverwash, unglaciated northern Great Plains type. Midwest Regional Office, Minneapolis, MN.

Thilenius, J.F. 1971. Vascular plants of the Black Hills of South Dakota and adjacent Wyoming. Research Paper RM-171. USDA Forest Service.